

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 39/395, 48/00, C12Q 1/02, 1/68,</b> <b>C12N 5/00, 15/00</b>	<b>A1</b>	<b>(11) International Publication Number: WO 97/36615</b> <b>(43) International Publication Date: 9 October 1997 (09.10.97)</b>
<b>(21) International Application Number:</b> PCT/US97/05440 <b>(22) International Filing Date:</b> 1 April 1997 (01.04.97)  <b>(30) Priority Data:</b> 60/014,570 1 April 1996 (01.04.96) US  <b>(71) Applicant:</b> PRESIDENT AND FELLOWS OF HARVARD COLLEGE [US/US]; 17 Quincy Street, Cambridge, MA 02138 (US).  <b>(72) Inventors:</b> PERRELLA, Mark, A.; 33 Pond Avenue #420, Brookline, MA 02146 (US). LEE, Mu-En; 102 Nardell Road, Newton, MA 02159 (US).  <b>(74) Agent:</b> FRASER, Janis, K.; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).		<b>(81) Designated States:</b> CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> REGULATION OF VASCULAR SMOOTH MUSCLE CELL HEME OXYGENASE-1  <b>(57) Abstract</b>  The invention features a method of inhibiting sepsis-associated hypotension in a mammal which includes the steps of identifying a mammal suffering from or at risk of developing sepsis and administering to the mammal a compound which inhibits expression of inducible heme oxygenase (HO-1). Also disclosed are screening assays useful for identifying a compound capable of inhibiting HO-1 expression or enzymatic activity.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

REGULATION OF VASCULAR SMOOTH MUSCLE CELL HEME  
OXYGENASE-1

Background of the Invention

5           The field of invention is treatment for  
hypotension. Septic shock, the most common cause of  
death in intensive care units, is characterized by severe  
and often irreversible hypotension (Parrillo, J. E.,  
1993, *N. Engl. J. Med.* 328:1471-1477). Sepsis leading to  
10 shock may be a detrimental consequence of a severe gram  
negative bacterial infection. Endotoxic shock is  
initiated by the release of bacterial cell wall-derived  
lipopolysaccharide (LPS, also known as endotoxin) and the  
subsequent production of cytokines and vasoactive  
15 mediators that result in vascular smooth muscle cell  
relaxation and hypotension (Glauser et al., 1991, *Lancet*  
338:732-736).

Summary of the Invention

          It has now been discovered that inducible heme  
20 oxygenase (HO-1) transcription and enzymatic activity are  
markedly increased in response to LPS, suggesting that  
HO-1 generated carbon monoxide (CO) contributes to the  
reduction in vascular tone during sepsis. Inhibition of  
sepsis-induced hypotension can be achieved by contacting  
25 the vascular cells with a compound that inhibits HO-1  
transcription and/or enzymatic activity.

          The invention features a method of inhibiting  
sepsis-associated hypotension in a mammal, e.g., a human  
patient, which includes the steps of identifying a mammal  
30 at risk of developing sepsis and administering to the  
mammal a compound which inhibits expression of HO-1. A  
mammal, such as a rat, mouse, rabbit, guinea pig,  
hamster, cow, pig, horse, goat, sheep, dog, cat, or  
human, at risk of developing sepsis is one that has

- 2 -

severe inflammation, e.g., due to infection by a bacterial, fungal, or viral pathogen. For example, hypotension associated with sepsis may be induced by a component of gram negative bacterial pathogens such as endotoxin, or by components produced by gram positive bacterial pathogens, viruses, and fungi, e.g., exotoxins which are secreted by such pathogens. Hypotension may also be induced by other inflammation-associated factors, such as cytokines, e.g., interleukin 1- $\beta$  (IL1- $\beta$ ) or tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ).

Preferably, the compound selectively inhibits HO-1 expression or activity, i.e., it inhibits expression of HO-1 without significantly inhibiting expression of non-inducible heme-oxygenase (HO-2). For example, the compound may inhibit HO-1 transcription in a mammalian vascular cell such as an aortic smooth muscle cell or endothelial cell by reducing HO-1 promoter activity, e.g., by blocking the binding of trans-acting transcription factors to HO-1 promoter sequences. Alternatively, the compound may inhibit HO-1 expression by inhibiting translation of HO-1 mRNA. For example, antisense therapy may be carried out by administering a single stranded nucleic acid complementary at least a portion of HO-1 mRNA to interfere with the translation of mRNA into protein, thus reducing the amount of functional HO-1 produced by the cell.

Also within the invention is a method of inhibiting sepsis-associated hypotension in a mammal which includes the steps of identifying a mammal at risk of developing sepsis and administering to the mammal a compound which inhibits the enzymatic activity of HO-1. The enzymatic activity of HO-1 is defined as the ability to generate carbon monoxide and/or bilirubin by catabolism of heme. For example, an antibody which binds to at or near the active site of HO-1 may be administered to reduce the enzymatic activity of HO-1.

- 3 -

Sepsis-associated hypotension may be diagnosed in vivo by administering to a patient an HO-1 specific antibody linked to a detectable label and imaging where the label localizes in the patient. An elevated level of label in the vascular tissue of the patient compared to a normal control level indicates that the patient may be at risk of developing or is suffering from sepsis-associated hypotension.

The invention also includes screening assays to identify compounds capable of inhibiting HO-1 transcription and/or HO-1 enzymatic activity. For example, a method for determining the ability of a candidate compound to inhibit HO-1 promoter activity may include the steps of (a) providing DNA comprising a HO-1 promoter sequence linked to a reporter gene, e.g., a gene encoding a detectable marker, e.g., luciferase or alkaline phosphatase; (b) contacting the DNA with a candidate compound; and (c) determining the level of expression of the reporter gene. A decrease in the level of expression in the presence of the compound, compared to the level of expression in the absence of the compound, is an indication of the ability of the candidate compound to inhibit HO-1 promoter activity. In another example, a method for determining the ability of a candidate compound to inhibit expression of HO-1 in vascular smooth muscle cells may include the steps of (a) providing a vascular smooth muscle cell; (b) contacting the cell with a candidate compound; and (c) determining the level of HO-1 expression in the cell. The method may also include the step of determining the amount of HO-2 in the vascular smooth muscle cell or a step of determining the amount of inducible nitric oxide synthase (iNOS) in the vascular smooth muscle cell. A decrease in the amount of HO-1 expression in the presence of the candidate compound compared to the amount in the absence of the candidate compound indicates that the

- 4 -

candidate compound inhibits expression of HO-1 in vascular smooth muscle cells. A larger decrease in the amount of HO-1 expression than in HO-2 expression indicates that the candidate compound selectively  
5 inhibits HO-1 expression. A concomitant decrease in the amount of iNOS expression indicates that the candidate compound inhibits expression of both HO-1 and iNOS.

The invention also includes a transgenic non-human mammal, e.g., a rodent such as a mouse, the germ cells  
10 and somatic cells of which contain a null mutation, e.g., a deletion, in DNA encoding HO-1. By "null mutation" is meant an alteration in the nucleotide sequence that renders the gene incapable of expressing a functional protein product. The mutation could be in the HO-1  
15 regulatory regions or in the coding sequence. It can, e.g., introduce a stop codon that results in production of a truncated, inactive gene product or it can be a deletion of all or a substantial portion of the coding sequence. For example, an exon, e.g., exon 3, of an HO-1  
20 gene may be deleted. The invention also includes a mammalian cell line, e.g., immortalized HO-1 deficient vascular smooth muscle cells, the genomic DNA of which contains a null mutation in DNA encoding HO-1. Such cells lack the ability to synthesize enzymatically-active  
25 HO-1. The cells harboring the null mutation may be derived from a vascular smooth muscle cell obtained from a HO-1 deficient transgenic mammal.

Other features and advantages of the invention will be apparent from the following detailed description,  
30 and from the claims.

#### Brief Description of the Drawings

Fig. 1 is a bar graph showing induction of heme oxygenase (HO) enzymatic activity by LPS in aortic tissue in vivo. Conscious male Sprague-Dawley rats (200-250 g)  
35 were injected with *Salmonella typhosa* LPS (n=2) at a dose of 4 mg/kg intravenously (i.v.) or vehicle as a control

- 5 -

(n=2). The rats were killed 9 hr after LPS or vehicle administration, and HO enzymatic activity was assessed in the aortic tissue. The experiment shown was performed in duplicate. HO enzymatic activity is expressed as a mean  
5  $\pm$  SD of the two experiments.

Fig. 2 is a bar graph showing a time-course of HO-1 mRNA induction by IL-1 $\beta$  in RASMC. Rat aortic smooth muscle cells (RASMC) were exposed to IL-1 $\beta$  (10 ng/ml), and total RNA was extracted from the cells at the  
10 indicated times. RNA was also extracted from cells receiving a vehicle (control; no IL-1 $\beta$ ) at the indicated times. Northern blot analyses were performed using 5  $\mu$ g of total RNA per lane. After electrophoresis, the RNA was transferred to nitrocellulose filters which were  
15 hybridized with a  $^{32}$ P-labeled rat HO-1 probe. The filters were also hybridized with a  $^{32}$ P-labeled oligonucleotide probe complementary to 18S ribosomal RNA to assess loading differences. The signal intensity of each RNA sample hybridized to the HO-1 probe was divided by that  
20 hybridized to the 18S probe. The normalized intensities were then plotted as a percentage of vehicle.

Fig. 3 is a bar graph showing a time course of HO-2 mRNA induction by IL-1 $\beta$  in RASMC. Filters from the experiment shown in Fig. 2 were rehybridized with a  
25  $^{32}$ P-labeled rat HO-2 probe. The signal intensity of each RNA sample hybridized to the HO-2 probe was divided by that hybridized to the 18S probe, and the normalized intensities were then plotted as a percentage of vehicle.

Fig. 4 is a bar graph showing dose-dependent HO-1  
30 mRNA induction by IL-1 $\beta$  in RASMC. RASMC were treated for 8 hours with the indicated concentrations of IL-1 $\beta$ . Total RNA was extracted at the end of the incubation period. Northern blot analyses were performed using 5  $\mu$ g of total RNA per lane. After electrophoresis, the RNA  
35 was transferred to nitrocellulose filters, which were hybridized to a  $^{32}$ P-labeled rat HO-1 probe. The filters

- 6 -

were also hybridized with a  $^{32}\text{P}$ -labeled oligonucleotide probe complementary to 18S ribosomal RNA to assess loading differences. The signal intensity of each RNA sample hybridized to HO-1 was divided by that hybridized to the 18S probe, and the normalized intensities were then plotted as a percentage of dose 0 ng/ml.

Fig. 5 is a bar graph showing the effect of IL- $1\beta$  on HO-1 mRNA transcription. Confluent RASMC were either stimulated with IL- $1\beta$  or vehicle for 24 hours. Nuclei were then isolated from the RASMC, and *in vitro* transcription was allowed to resume in the presence of [ $\alpha$ - $^{32}\text{P}$ ] UTP. Equal amounts of  $^{32}\text{P}$ -labeled, *in vitro* transcribed RNA probes from each group were hybridized to 1  $\mu\text{g}$  of denatured HO-1,  $\beta$ -actin, and HO-2 cDNA that had been immobilized on nitrocellulose filters. The signal intensity of HO-1 was normalized by reference to the  $\beta$ -actin signal, and the transcriptional rate was plotted as a percentage of vehicle. Fig. 6 is a line graph showing stability of HO-1 message, as determined by measuring mRNA levels in the presence of the transcription inhibitor actinomycin D. RASMC were stimulated with IL- $1\beta$  (10 ng/ml) for 12 hours, followed by administration of actinomycin D (5  $\mu\text{g}/\text{ml}$ ). Total RNA was extracted from the RASMC at the indicated times after administration of actinomycin D. Northern blot analyses were performed using 5  $\mu\text{g}$  of total RNA per lane. After electrophoresis, the RNA was transferred to nitrocellulose filters, which were hybridized to  $^{32}\text{P}$ -labeled HO-1 and 18S probes. To correct for differences in loading, the signal intensity of each RNA sample hybridized to the HO-1 probe was divided by that hybridized to the 18S probe. The normalized intensity was then plotted as a percentage of the 0-hour value against time (in log scale).

Fig. 7 is a bar graph showing the effect of cycloheximide (CHX) on HO-1 mRNA induction by IL- $1\beta$  in



- 7 -

RASMC. RASMC were exposed to no stimulus (control), IL-1 $\beta$  (10 ng/ml), CHX alone (10  $\mu$ g/ml, CHX control), or a combination of CHX (10  $\mu$ g/ml) and IL-1 $\beta$  (10 ng/ml). CHX was applied to the cells one-half hour before IL-1 $\beta$  was applied. Total RNA was extracted from the cells after 8 hours of IL-1 $\beta$  stimulation, and Northern blot analysis was performed using 5  $\mu$ g of total RNA per lane. After electrophoresis the RNA was transferred to nitrocellulose filters, which were hybridized to <sup>32</sup>P-labeled HO-1 and 18S probes. The signal intensity of each RNA sample hybridized to HO-1 was divided by that hybridized to the 18S probe, and the normalized intensities were then plotted as a percentage of control. The bars represent mean  $\pm$  SE of three separate experiments.

15

#### Detailed Description

HO generates biliverdin (which is subsequently reduced to bilirubin) and CO, a potent vasodilator that contributes to the reduction in vascular tone and hypotension associated with sepsis. HO-1 enzymatic activity was found to be upregulated within vascular tissue by a pathophysiologic process, sepsis, *in vivo*. LPS was found to markedly increase both HO-1 mRNA levels and HO enzymatic activity in vascular tissue *in vivo*, and IL-1 $\beta$  was found to induce HO-1 mRNA in rat aortic smooth muscle cells (RASMC) *in vitro*. In both large blood vessels (aorta) and small resistance vessels (arterioles), the increase in staining for HO-1 localized to vascular smooth muscle cells and endothelial cells. The induction of vascular smooth muscle cell-derived HO-1 *in vitro* occurred at the level of gene transcription. The antioxidants pyrrolidinedithiocarbamate (PDTC) and N-acetylcysteine (NAC) did not prevent the induction of HO-1 mRNA by IL-1 $\beta$ , suggesting this induction was not the result of oxidative stress.

- 8 -

The marked induction of HO-1 enzymatic activity within vascular tissue by LPS suggests that the CO generated by HO enzymatic activity contributes to the reduction in vascular tone during endotoxic shock.

5 Agents which selectively inhibit HO-1 transcription and/or those which selectively inhibit HO-1 enzymatic activity, e.g., selective antagonists of HO-1, can be administered to patients to prevent and treat sepsis-associated hypotension. The role of CO in endotoxic  
10 shock was evaluated by analyzing the regulation of vascular HO-1 in vivo and investigating the mechanism of HO-1 regulation in vascular smooth muscle cells in vitro as described below.

#### Cell culture

15 RASMC were harvested from male Sprague-Dawley rats (200-250 grams) by enzymatic dissociation using standard methods. The cells were cultured in Dulbecco's modified Eagle's medium (DMEM, JRH Biosciences, Lenexa, KS) and supplemented with 10% fetal calf serum, penicillin  
20 (100 U/ml), streptomycin (100 µg/ml), and 25 mM Herpes (pH 7.4) (Sigma Chemical, St. Louis, MO). RASMC were passaged in culture every 4 to 7 days, and experiments were performed on cells which were 4 to 6 passages from primary culture. After the cells had grown to  
25 confluence, they were placed in 2% fetal calf serum 12 hours before commencement of the experiments. *Salmonella typhosa* LPS (Sigma Chemical) was dissolved in phosphate buffered saline and stored at -20°C. Recombinant human IL-1β (Collaborative Biomedical, Bedford, MA) was stored  
30 at -80°C until use. PDTC (Sigma Chemical) was dissolved in sterile water and stored at 4°C, and NAC (L form, Chiron Therapeutics, Emeryville, CA) was diluted in sterile water and used immediately.

RNA blot hybridization

Total RNA was obtained from rat aortas and cultured smooth muscle cells by guanidinium isothiocyanate extraction and centrifuged through cesium chloride. The RNA was fractionated on a 1.3% formaldehyde-agarose gel and transferred to nitrocellulose filters. The filters were hybridized at 68°C for 2 hr with <sup>32</sup>P-labeled rat HO-1 or HO-2 probes in QuikHyb solution (Stratagene, La Jolla, CA). The hybridized filters were then washed in 30 mM sodium chloride, 3 mM sodium citrate, and 0.1% sodium dodecyl sulfate solution at 55°C and autoradiographed with Kodak XAR film at -80°C for 4 to 12 hours or stored on phosphor screens for 2 to 4 hours. To correct for differences in RNA loading, the filters were washed in a 50% formamide solution at 80°C and rehybridized with an 18S ribosomal RNA probe. Images were displayed and radioactivity was measured on a PhosphorImager running the ImageQuant software (Molecular Dynamics, Sunnyvale, CA).

Induction of HO-1 mRNA by IL-1 $\beta$  in response to the antioxidants PDTC and NAC was measured as follows. For PDTC experiments, RASMC were exposed to no stimulus (control), IL-1 $\beta$  (10 ng/ml), IL-1 $\beta$  (10 ng/ml) plus PDTC (25 and 50  $\mu$ M), or PDTC alone (25 and 50  $\mu$ M). For NAC experiments, RASMC were exposed to no stimulus (control), IL-1 $\beta$  (10 ng/ml), IL-1 $\beta$  (10 ng/ml) plus NAC (5 and 25 mM), or NAC alone (5 mM). Twenty-four hours after stimulation, total RNA was extracted from the cells. Northern blot analyses were performed using 10  $\mu$ g of total RNA per lane. After electrophoresis, the RNA was transferred to nitrocellulose filters, which were hybridized to <sup>32</sup>P-labeled HO-1 and 18S probes.

Nuclear run-on analysis

RASMC were either not stimulated (vehicle) or stimulated with IL-1 $\beta$  (10 ng/ml) for 24 hours. The cells were subsequently lysed and nuclei were isolated as

- 10 -

described using standard protocols. Nuclear suspension (200  $\mu$ l) was incubated with 0.5 mM each of CTP, ATP, and GTP and with 125  $\mu$ Ci of  $^{32}$ P-labeled UTP (3,000 Ci/mmol) (Du Pont-New England Nuclear). The samples were  
5 extracted with phenol/chloroform, precipitated, and resuspended at equal counts/min/ml in hybridization buffer ( $9.7 \times 10^6$  cpm/ml). Denatured probes (1  $\mu$ g) dot-blotted on nitrocellulose filters were hybridized at 40°C for four days in the presence of formamide. HO-1  
10 and HO-2 cDNA fragments were amplified from RASMC RNA by the reverse transcription polymerase chain reaction (PCR) (Morita et al., 1995, Proc. Natl. Acad. Sci. USA 92:1475-1479). HO-1 and HO-2 cDNA were used as probes to detect the presence of HO-1 and HO-2 transcripts, and  $\beta$ -actin  
15 cDNA was used as a control probe.

#### HO enzymatic activity

Aortas were harvested from rats treated with vehicle or LPS (4 mg/kg i.v.), the adventitia was stripped, and the tissue was homogenized in buffer  
20 containing 30 mM Tris, pH 7.5, 0.25 M sucrose, 0.15 M NaCl, and the Complete™ protease inhibitor (Boehringer Mannheim) using a Polytron™ homogenizer. The homogenate was filtered through two layers of cheesecloth, and centrifuged at 10,000  $\times$  g for 15 min. The supernatant  
25 fraction was subsequently centrifuged at 100,000  $\times$  g for 1 h. The microsomal pellet was resuspended in 50 mM potassium phosphate buffer (pH 7.4) containing Complete™ protease inhibitor. Heme oxygenase activity was measured by bilirubin generation using standard methods. The  
30 liver microsomal supernatant fraction from the control animal served as the source of biliverdin reductase. A reaction mixture (0.5 ml) containing 33  $\mu$ M hemin, rat liver microsomal supernatant fraction (0.3 mg), NADPH generating system, and aortic microsomal protein was  
35 incubated at 37°C for 10 min in the dark. The reaction mixture without NADPH generating system was used as a

- 11 -

control. The reactions were stopped by placement on ice, and subsequently scanned with a spectrophotometer (Beckman, Columbia, MD). The amount of bilirubin formed was determined as the difference in optical density units  
5 between 462 nm and 530 nm (extinction coefficient,  $40 \text{ nm}^{-1}\text{cm}^{-1}$  for bilirubin). HO enzymatic activity was expressed as nmol of bilirubin formed per mg of protein per hour. The protein concentration was determined by a dye-binding assay (Bio-Rad, Hercules, CA).

#### 10 Immunocytochemical staining

Adult male Sprague-Dawley rats treated with LPS or vehicle were perfused with 4% paraformaldehyde. The aortas were removed, post-fixed with 4% paraformaldehyde overnight at 4°C, and then soaked in 30% sucrose for 2  
15 days at 4°C. The specimens were cut at a thickness of 5  $\mu\text{m}$ . Immunocytochemical staining of tissue sections was performed using standard methods. To reduce nonspecific binding, the sections were incubated in phosphate buffered saline containing 10% normal goat serum and 0.4%  
20 triton X-100 for 30 minutes. Rabbit polyclonal antibody against purified rat liver HO-1 (StressGen Biotechnologies, Victoria, BC, Canada) was applied for 1 hour at room temperature and then overnight at 4°C at a dilution of 1:1000-1200. Sections were rinsed twice with  
25 high-salt phosphate buffered saline (0.5 M NaCl) and once with regular phosphate buffered saline (5 minutes for each wash) and then incubated with biotinylated goat anti-rabbit IgG (Vector Labs, Burlingame, CA) at a dilution of 1:500 for 1 hour at room temperature. The  
30 sections were then rinsed with phosphate buffered saline and incubated with avidin-biotin complex (ABC elite kit, Vector Labs, Burlingame, CA) at a dilution of 1:100 for 1 hour at room temperature. After washing with PBS, the tissue sections were treated with diaminobenzidine (DAB)  
35 in phosphate buffered saline- $\text{H}_2\text{O}_2$  for 1-3 minutes using the peroxidase substrate kit DAB (Vector Labs,

- 12 -

Burlingame, CA) and then transferred into phosphate-buffered saline solution to stop the reaction. The presence of HO-1 was indicated by the development of a brown color within the cytoplasm. Counterstaining was performed with 0.5% methyl green.

For immunocytochemical localization of HO-1 expression in vascular tissue, conscious male Sprague-Dawley rats (200-250 grams) were injected with vehicle or *Salmonella typhosa* LPS at a dose of 4 mg/kg i.v. The rats were killed 9 hr after vehicle or LPS administration and perfused with 4% paraformaldehyde. The aortic tissue was removed and immunocytochemical staining was performed as described above. Stained aortic tissue was viewed at x190 magnification, and stained adventitial arterioles (vessels less than 300  $\mu$ m in diameter) were viewed at x540 magnification.

#### Data analysis

Comparisons between the vehicle and LPS-treated groups for HO enzymatic activity were made using unpaired t-tests (two tailed). Statistical significance was accepted for a p value <0.05.

#### LPS induces vascular HO-1 mRNA and enzymatic activity in vivo

To determine if LPS regulates vascular HO-1 in an animal model of sepsis, rats were injected with vehicle or *Salmonella typhosa* LPS (4 mg/kg i.v.) as described above. HO-1 mRNA levels were found to be markedly increased in aortic tissue after 9 hours of LPS stimulation compared to the levels in tissue from rats receiving vehicle (Fig. 1). The administered dose of LPS produces hypotension in rats; the 9 hour time point at which HO-1 transcription was evaluated was chosen after performing an *in vivo* time course experiment. LPS was found not to induce an increase in HO-2 mRNA levels.

The following experiments were carried out to determine whether the increase in HO-1 mRNA levels

- 13 -

corresponded to an increase in HO enzymatic activity. Rats were given vehicle or LPS (4 mg/kg i.v.), and the aortas were harvested 9 hours later (adventitia of the vessels was stripped prior to analysis). LPS promoted an 8.9-fold increase ( $p < 0.05$ ) in HO enzymatic activity (Fig. 1). In fact, the level of HO enzymatic activity in the aortic tissue from rats receiving LPS ( $36.3 \pm 2.4$  nmol/mg protein/hour) was comparable to the level of HO activity in the liver of control rats ( $40 \pm 2.1$  nmol/mg protein/hour,  $p = 0.34$ ). These data indicate that a significant amount of inducible HO enzymatic activity is generated in vascular tissue after LPS stimulation.

To localize the arterial cell type within the vessel responsible for the increase in HO enzymatic activity, immunocytochemical staining was performed using a rabbit anti-HO-1 antibody. Staining for HO-1 protein was increased in the smooth muscle cells of aortas from rats receiving LPS compared with rats receiving vehicle. Immunocytochemical staining demonstrated an LPS-induced increase in HO-1 expression in the smooth muscle cells of arterioles (smaller vessels that contribute to the regulation of vascular tone). Staining for HO-1 was also increased in the endothelium of aortas and arterioles after LPS stimulation.

These data indicate that HO enzymatic activity, vascular HO-1 mRNA levels, and protein levels are increased after LPS administration *in vivo*.  
IL-1 $\beta$  induces HO-1, but not HO-2, mRNA in vascular smooth muscle cells *in vitro*

Northern blot analyses were performed with total RNA from RASMC exposed to either vehicle (control) or IL-1 $\beta$ . The blots were then hybridized to HO-1 and HO-2 cDNA probes. The time course (0 to 48 hours) of HO-1 mRNA stimulation by IL-1 $\beta$  (10 ng/ml) is shown in Fig. 2. IL-1 $\beta$  induced HO-1 mRNA as early as 4 hours after stimulation, and peak induction occurred after 24 hours.

Twenty four hours after the administration of IL-1 $\beta$ , HO-1 mRNA increased by 5.8-fold compared to the control. The induction of HO-1 mRNA by IL-1 $\beta$  decreased to 2.8-fold after 48 hours. The message for HO-2, in contrast to HO-1, increased only minimally after treatment with IL-1 $\beta$  (Fig. 3).

RASMC were also stimulated for 8 hours with increasing doses of IL-1 $\beta$  (Fig. 4). Maximal induction of HO-1 mRNA occurred at a dose of 10 ng/ml, although as little as 0.1 ng/ml promoted an increase in mRNA levels. Increasing the dose of IL-1 $\beta$  beyond 10 ng/ml did not further increase HO-1 mRNA levels significantly.

#### IL-1 $\beta$ increases the rate of HO-1 gene transcription

To determine the mechanism by which IL-1 $\beta$  increases HO-1 mRNA in vascular smooth muscle cells, experiments to assess the transcriptional rate and stability of HO-1 mRNA were performed. Nuclear run-on experiments showed a 5-fold increase in the rate of HO-1 gene transcription after 24 hours of IL-1 $\beta$  stimulation. Stability of HO-1 mRNA transcripts was determined by measuring mRNA levels in the presence of the transcription inhibitor actinomycin D. HO-1 mRNA was found to have a half-life of 1.3 hours after exposure to vehicle (Fig. 6). The half-life of HO-1 mRNA was not prolonged by IL-1 $\beta$ . These data indicate that the increase in HO-1 mRNA levels by IL-1 $\beta$  stimulation is the result of an increase in HO-1 gene transcription.

#### Induction of HO-1 mRNA by IL-1 $\beta$ in RASMC requires protein synthesis

RASMC were treated with the protein synthesis inhibitor cycloheximide (10  $\mu$ g/ml) for 1 hour before adding IL-1 $\beta$  (10 ng/ml). This dose of cycloheximide completely inhibits leucine uptake in RASMC. Cycloheximide completely inhibited the induction of HO-1 mRNA by IL-1 $\beta$  at 8 hours (Fig. 7). This study was also performed using another protein synthesis inhibitor,



- 15 -

anisomycin. Anisomycin also prevented IL-1 $\beta$ -induced HO-1 mRNA induction. These data indicate that new protein synthesis is required for the induction of HO-1 mRNA by IL-1 $\beta$  in vascular smooth muscle cells.

5 Induction of HO-1 mRNA by IL-1 $\beta$  is not prevented by antioxidants

To determine if oxidative stress promotes the induction of HO-1 mRNA by IL-1 $\beta$ , RASMC were treated with IL-1 $\beta$  (10 ng/ml) for 24 hours in the presence or absence  
10 of the antioxidant PDTC (25 and 50  $\mu$ M). PDTC did not inhibit IL-1 $\beta$ -induced HO-1 mRNA induction. This study was repeated using another antioxidant, N-acetylcysteine (NAC). NAC (5 and 25 mM) also had no effect on the induction of HO-1 mRNA by IL-1 $\beta$ . Previous studies have  
15 shown that comparable doses of PDTC and NAC inhibit the expression of genes upregulated by IL-1 $\beta$ -induced oxidative stress (Marui et al., 1993, J. Clin. Invest. 92:1866-1874). However, these data indicate that oxidative stress did not contribute to the induction of  
20 HO-1 mRNA by IL-1 $\beta$  in vascular smooth muscle cells.

HO-1 deficient transgenic animals

An HO-1 deficient transgenic animal, e.g., a HO-1 knockout mouse, can be produced as follows.

The murine HO-1 gene contains 5 exons and 4  
25 introns, spanning approximately 7 kilobases (kb). The targeting construct was made by deleting the largest exon (exon 3) which contains 492 nucleotides out of the 867 nucleotides of the entire open reading frame. This deletion renders the HO-1 enzyme non-functional. A rat  
30 HO-1 cDNA was used to isolate genomic clones from a 129/SV mouse genomic library. The XhoI/BamHI fragment of the neo cassette from pMC1neo PolyA plasmid was subcloned into pBluescript II SK (Stratagene, La Jolla, CA) to generate pBS-neo. To generate pBS-neo-HO-1, the 3 kb  
35 XhoI fragment of the HO-1 gene spanning from exon 1 to the end of intron 2 was subcloned into the XhoI site of

pBS-neo in the same orientation as the neo cassette. The 4 kb HO-1 *Bam*HI-*Eco*RI fragment containing a small portion of intron 3, exon 4, and exon 5 was subcloned into *Bam*HI and *Eco*RI site of pPGK-TK to generate pPGK-TK-HO-1. The 7 kb *Bam*HI-*Cla*I fragment (filled in with Klenow) from pPGK-TK-HO-1 was then subcloned into *Bam*HI and *Xba*I sites (filled in with Klenow) sites of pBS-neo-HO-1 to generate the HO-1 targeting construct.

The linearized targeting construct is transfected into murine D3 embryonic stem (ES) cells, and a clone with the correct homologous recombination (yielding the appropriately disrupted HO-1 gene) injected into blastocysts and used to generate HO-1 deficient mice. HO-1 deficient mice and HO-1 deficient cell lines are useful in determining the etiology of sepsis-associated hypotension.

#### Therapeutic and diagnostic uses

The marked increase in HO-1 transcription and enzymatic activity suggests that HO-1-generated CO contributes to the reduction in vascular tone during endotoxic shock. Inhibition of sepsis-induced hypotension can be achieved by contacting the vascular cells with a compound that inhibits HO-1 transcription and/or enzymatic activity

Prior to the invention, it was thought that further induction of HO-1 by hemoglobin may be of therapeutic benefit in endotoxic shock (Otterbein et al., 1995, Am. J. Respir. Cell Mol. Biol. 13:595-601) because the generation of bilirubin (which has antioxidant properties) would protect against endotoxin-induced oxidative damage in the lung. The data described herein indicates that CO production (the other product of heme metabolism), as a result of the marked induction of vascular HO-1 enzymatic activity, contributes to the

- 17 -

decline in vascular tone during endotoxic shock and thus worsens the patient's condition.

Nucleic acids complementary to all or part of the HO-1 coding sequence (GenBank Accession No. X06985; Yoshida et al., 1988, Eur. J. Biochem. 171:457-461 (TABLE 1)) may be used in methods for antisense treatment to inhibit expression of HO-1. Antisense treatment may be carried out by administering to a mammal, such as a human, DNA containing a promoter, e.g., a vascular smooth muscle cell-specific promoter or an endothelial cell-specific promoter, operably linked to a DNA sequence (an antisense template), which is transcribed into an antisense RNA. Alternatively, as mentioned above, antisense oligonucleotides may be introduced directly into vascular smooth muscle cells. The antisense oligonucleotide may be a short nucleotide sequence (generally at least 10, preferably at least 14, more preferably at least 20 (e.g., at least 30), and up to 100 or more nucleotides) formulated to be complementary to a portion, e.g., the coding sequence, or all of HO-1 mRNA. Standard methods relating to antisense technology have been described (see, e.g., Melani et al., 1991, Cancer Res. 51:2897-2901). Following transcription of a DNA sequence into an antisense RNA, the antisense RNA binds to its target nucleic acid molecule, such as an mRNA molecule, thereby inhibiting expression of the target nucleic acid molecule. For example, an antisense sequence complementary to a portion or all of HO-1 mRNA could be used to inhibit the expression of HO-1, thereby decreasing sepsis-associated hypotension in a mammal. Oligonucleotides complementary to various sections of HO-1 mRNA can readily be tested *in vitro* for their ability to decrease production of HO-1, using assays similar to those described herein. Promising oligonucleotides can then be tested *in vivo* in rats or mice challenged with LPS.

Any vector that can transfect a vascular smooth muscle cell may be used in this method of the invention. Preferred vectors are viral vectors, including those derived from replication-defective hepatitis viruses (e.g., HBV and HCV), retroviruses (see, e.g., WO 89/07136; Rosenberg et al., 1990, N. Eng. J. Med. 323(9):570-578), adenovirus (see, e.g., Morsey et al., 1993, J. Cell. Biochem., Supp. 17E), adeno-associated virus (Kotin et al., 1990, Proc. Natl. Acad. Sci. USA 87:2211-2215), replication defective herpes simplex viruses (HSV; Lu et al., 1992, Abstract, page 66, Abstracts of the Meeting on Gene Therapy, Sept. 22-26, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York), and any modified versions of these vectors. The invention may utilize any other delivery system which accomplishes *in vivo* transfer of nucleic acids into eukaryotic cells. For example, the nucleic acids may be packaged into liposomes, receptor-mediated delivery systems, non-viral nucleic acid-based vectors, erythrocyte ghosts, or microspheres (e.g., microparticles); see, e.g., U.S. Patent No. 4,789,734; U.S. Patent No. 4,925,673; U.S. Patent No. 3,625,214; Gregoriadis, 1979, Drug Carriers in Biology and Medicine, pp. 287-341 (Academic Press,). Alternatively, naked DNA may be administered. Delivery of nucleic acids to a specific site in the body for antisense therapy may also be accomplished using a biolistic delivery system, such as that described by Williams et al., 1991, Proc. Natl. Acad. Sci. USA 88:2726-2729.

Antisense oligonucleotides may consist of DNA, RNA, or any modifications or combinations thereof. As an example of the modifications that the oligonucleotides may contain, inter-nucleotide linkages other than phosphodiester bonds, such as phosphorothioate, methylphosphonate, methylphosphodiester, phosphorodithioate, phosphoramidate, phosphotriester, or

- 19 -

phosphate ester linkages (Uhlman et al., 1990, Chem. Rev. 90(4):544-584; Anticancer Research, 1990, 10:1169) may be present in the oligonucleotides, resulting in their increased stability. Oligonucleotide stability may also  
5 be increased by incorporating 3'-deoxythymidine or 2'-substituted nucleotides (substituted with, e.g., alkyl groups) into the oligonucleotides during synthesis, by providing the oligonucleotides as phenylisourea derivatives, or by having other molecules, such as  
10 aminoacridine or poly-lysine, linked to the 3' ends of the oligonucleotides (see,

- 20 -

TABLE 1

Human HO-1 cDNA

```

1   tcaacgcctg cctccccctg agcgtcctca ggcagccgc cgcgcgcgga gccagcacga
61  acgagcccag caccggcccg atggagcgtc cgcaaccga cagcatgccc caggatttgt
5  121 cagaggccct gaaggaggcc accaaggagg tgcacacca ggcagagaat gctgagttca
181 tgaggaactt tcagaagggc caggtgacct gagacggctt caagctggtg atggcctccc
241 tgtaccacat ctatgtggcc ctggaggagg agattgagcg caacaaggag agcccagttc
301 tcgcccctgt ctacttccca gaagagctgc accgcaaggc tgccctggag caggacctgg
361 ccttctggtg cgggccccgc tggcaggagg tcatccccta cacaccagcc atgcagcgct
10 421 atgtgaagcg gctccacgag gtggggcgca cagagcccga gctgctggtg gcccacgect
481 acaccgcta cctgggtgac ctgtctgggg gccaggtgct caaaaagatt gcccgaaag
541 ccctggacct gccagctctt ggcgagggcc tggccttctt caccttcccc aacattgcc
601 gtgccaccaa gttcaagcag ctctaccgct cccgcatgaa ctccctggag atgactcccg
661 cagtcaggca gaggggtgata gaagaggcca agactgcgtt cctgctcaac atccagctct
15 721 ttgaggagtt gcaggagctg ctgacctatg acaccaagga ccagagcccc tcacgggac
781 cagggtctcg ccagcgggcc agcaacaaag tgcaagattc tgcccccggt gagactccca
841 gagggagacc cccactcaac acccgctccc aggtccgctt tctccgatgg gtccttacac
901 tcagctttct ggtggcgaca gttgctgtag ggctttatgc catgtgaatg caggcatgct
961 ggctcccagg gccatgaact ttgtccggtg gaaggccttc tttctagaga gggaattctc
20 1021ttggctggct tccttaccgt gggcactgaa ggctttcagg gcctccagcc ctctcactgt
1081gtccctctct ctggaaagga ggaaggagcc tatggcatct tcccacga aaagcacatc
1141caggcaatgg cctaaacttc agagggggcg aaggggtcag ccctgccctt cagcatcctc
1201agttcctgca gcagagcctg gaagacaccc taatgtggca gctgtctcaa acctccaaaa
1261gccctgagtt tcaagtatcc ttgttgacac ggccatgacc actttccccg tgggcatgg
25 1321caatttttac acaaacctga aaagatgttg tgtcttgtgt ttttgttta ttttgtttg
1381agccactctg ttctggctc agcctcaaat gcagtatttt tgttgtgttc tgttgtttt
1441atagcagggg tgggggtggt tttgagccat gcgtgggtgg ggaggagggt gtttaacggc
1501actgtggcct tgggtctaact tttgtgtgaa ataataaaca acattgtctg

```

(SEQ ID NO:1)

e.g., Anticancer Research, 1990, 10:1169-1182).

Modifications of the RNA and/or DNA nucleotides may be present throughout the oligonucleotide, or in selected regions of the oligonucleotide, e.g., in the 5' and/or 3' ends. The antisense oligonucleotides may also be modified so as to increase their ability to penetrate the target tissue by, e.g., coupling the oligonucleotides to lipophilic compounds. Antisense oligonucleotides based on the HO-1 nucleotide sequence (TABLE 1) can be made by any method known in the art, including standard chemical synthesis, ligation of constituent oligonucleotides, and transcription of DNA complementary to the all or part of the HO-1 coding sequence.

HO-1 is naturally expressed and upregulated (in response to sepsis) in vascular smooth muscle cells and endothelial cells. These cells are, therefore, the preferred cellular targets for antisense therapy. Targeting of antisense oligonucleotides to vascular smooth muscle cells may be achieved, for example, by coupling the oligonucleotides to ligands of vascular smooth muscle cell receptors. Similarly, oligonucleotides may be targeted to vascular smooth muscle cells by being conjugated to monoclonal antibodies that specifically bind to vascular smooth muscle-specific cell surface proteins.

Therapeutic applications of antisense oligonucleotides in general are described, e.g., in the following review articles: Le Doan et al., Bull. Cancer 76:849-852, 1989; Dolnick, Biochem. Pharmacol. 40:671-675, 1990; Crooke, Annu. Rev. Pharmacol. Toxicol. 32, 329-376, 1992. Antisense nucleic acids may be used alone or in admixture, or in chemical combination, with one or more materials, including other antisense oligonucleotides or recombinant vectors, materials that increase the biological stability of the oligonucleotides or the recombinant vectors, or materials that increase

- 22 -

the ability of the therapeutic compositions to penetrate vascular smooth muscle cells selectively.

Therapeutic compositions, e.g., inhibitors of HO-1 transcription or enzymatic activity, may be administered  
5 in pharmaceutically acceptable carriers (e.g., physiological saline), which are selected on the basis of the mode and route of administration and standard pharmaceutical practice. Suitable pharmaceutical carriers, as well as pharmaceutical necessities for use  
10 in pharmaceutical formulations, are described in *Remington's Pharmaceutical Sciences*, a standard reference text in this field, and in the USP/NF. The compound may be administered with intravenous fluids as well as in combination with other anti-inflammatory agents, e.g.,  
15 antibiotics; glucocorticoids, such as dexamethasone (Dex). The compound may also be administered together with pressors, such as epinephrine or norepinephrine.

A therapeutically effective amount is an amount which is capable of producing a medically desirable  
20 result in a treated animal. As is well known in the medical arts, dosage for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health,  
25 and other drugs being administered concurrently. Dosages will vary, but a preferred dosage for intravenous administration of DNA is approximately  $10^6$  to  $10^{22}$  copies of the DNA molecule. The compositions of the invention may be administered locally or systemically.  
30 Administration will generally be parenterally, e.g., intravenously. As mentioned above, DNA may also be administered directly to the target site, e.g., by biolistic delivery to an internal or external target site or by catheter to a site in an artery.

35 Instead of inhibiting HO-1 transcription and/or translation, the enzymatic activity of HO-1 may be



inhibited to treat sepsis-associated hypotension in a mammal. For example, an antibody which binds to HO-1 may be administered to reduce the enzymatic activity of HO-1. For administration to human patients, antibodies, e.g.,  
5 HO-1 specific monoclonal antibodies, can be humanized by methods known in the art. Antibodies with a desired binding specificity can be commercially humanized (Scotgene, Scotland; Oxford Molecular, Palo Alto, CA).  
Identification and use of compounds which inhibit HO-1  
10 transcription or enzymatic activity

Compounds that inhibit HO-1 transcription and/or enzymatic activity can be identified by methods ranging from rational drug design to screening of random compounds. The latter method is preferable, as simple  
15 and rapid assays for testing such compounds are available. Small organic molecules are desirable candidate compounds for this analysis, as frequently these molecules are capable of passing through the plasma membrane to inhibit HO-1 transcription within the cell.

20 The screening of compounds for the ability to HO-1 transcription be carried by identifying compounds that block the binding of trans-acting factors to HO-1 promoter sequences. For example, AP-1 sites in the HO-1 promoter, e.g., TGTGTCA (SEQ ID NO:2) and/or TGAGTCA (SEQ  
25 ID NO:3) which may be targets of LPS-induced trans-acting factors in vascular smooth muscle cells are linked to a functional promoter and a reporter gene, e.g., the gene encoding luciferase or alkaline phosphatase. For luciferase constructs, the cells harboring the construct  
30 are harvested after exposure to the candidate compound and luciferase activity measured; for alkaline phosphatase constructs, the culture medium of the cells is collected and the amount of alkaline phosphatase secreted by the cells into the medium is measured.  
35 Stable promoter constructs containing both SEQ ID NO:2 and SEQ ID NO:3 and constructs lacking both sequences

- 24 -

have been made and introduced into vascular smooth muscle cells. Trans-acting factors which bind to SEQ ID NO:2 and/or SEQ ID NO:3 can be identified using these constructs.

5 To identify compounds capable of inhibiting HO-1 transcription, these cells containing SEQ ID NO:2 and/or SEQ ID NO:3 are contacted with candidate compounds and the ability the cells to generate the reporter protein is determined (e.g., luciferase in the cells or alkaline  
10 phosphatase in the media). A decrease in the amount of expression of the reporter protein indicates that the candidate compound inhibits HO-1 transcription.

Candidate compounds may also be screened using cell culture assays. Cells expressing HO-1, e.g.,  
15 vascular smooth muscle cells, are cultured in the presence of the candidate compound. The level of expression of HO-1 in the presence and absence of the compound may be measured using known methods, e.g., PCR or Northern blot analysis to measure transcription.  
20 Western blot analysis can be used to detect the presence of the HO-1 protein.

Alternatively, bilirubin or CO production may be measured to evaluate HO-1 enzymatic activity. To identify compounds capable of inhibiting HO-1 activity,  
25 vascular smooth muscle cells, e.g., cultured cells or primary cells derived from aortic tissue, are contacted with a candidate compound. A control sample of cells is processed in parallel in the absence of a candidate compound. HO-1 enzymatic activity is measured in both  
30 samples either by measuring bilirubin as described above or CO by gas chromatography. A decrease in the amount of bilirubin or CO in the presence of the candidate compound compared to that in the absence of the candidate compound indicates that the candidate compound inhibits HO-1  
35 enzymatic activity.

A decrease in the level of expression HO-1 indicates that the compound can inhibit sepsis-associated hypotension. Conversely, an increase in expression of HO-1 indicates that the compound can increase vascular hypotension. Such a compound can be used therapeutically to upregulate HO-1 expression, e.g., when administered locally in the lung, to treat pulmonary hypertension.

The therapeutic compounds identified using the methods of the invention may be administered to a patient by any appropriate method for the particular compound, e.g., orally, intravenously, parenterally, transdermally, transmucosally, by inhalation, or by surgery or implantation at or near the site where the effect of the compound is desired (e.g., with the compound being incorporated into a solid or semi-solid biologically compatible and resorbable matrix). Therapeutic doses are determined specifically for each compound, most being administered within the range of 0.001 to 100.0 mg/kg body weight, or within a range that is clinically determined to be appropriate by one skilled in the art.

In some cases, it may be advantageous to administer HO-1 inhibitors in combination with drugs that affect other points of a pathway leading to severe hypotension or septic shock. For example, an HO-1 inhibitor may be administered either simultaneously or sequentially with another therapeutic agent, e.g., TGF $\beta$  to inhibit iNOS or an antibiotic or anti-inflammatory drug to treat the underlying infection.

Other embodiments are within the claims.

- 26 -

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: The President and Fellows of Harvard College
- (ii) TITLE OF INVENTION: REGULATION OF VASCULAR SMOOTH MUSCLE  
CELL HEME OXYGENASE-1
- (iii) NUMBER OF SEQUENCES: 3
- (iv) CORRESPONDENCE ADDRESS:  
(A) ADDRESSEE: Fish & Richardson P.C.  
(B) STREET: 225 Franklin Street  
(C) CITY: Boston  
(D) STATE: MA  
(E) COUNTRY: USA  
(F) ZIP: 02110-2804
- (v) COMPUTER READABLE FORM:  
(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:  
(A) APPLICATION NUMBER:  
(B) FILING DATE:  
(C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:  
(A) NAME: Fraser, Janis K.  
(B) REGISTRATION NUMBER: 34,819  
(C) REFERENCE/DOCKET NUMBER: 05433/023W01
- (ix) TELECOMMUNICATION INFORMATION:  
(A) TELEPHONE: 617/542-5070  
(B) TELEFAX: 617/542-8906  
(C) TELEX: 200154

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 1550 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:
- |    |   |     |
|----|---|-----|
| 40 | TCAACGCCTG CCTCCCTCG AGCGTCCTCA GCGCAGCCGC CGCCCGCGGA GCCAGCACGA  | 60  |
|    | ACGAGCCCAG CACCGGCCGG ATGGAGCGTC CGCAACCCGA CAGCATGCCC CAGGATTTGT | 120 |
|    | CAGAGGCCCT GAAGGAGGCC ACCAAGGAGG TGCACACCCA GGCAGAGAAT GCTGAGTTCA | 180 |
|    | TGAGGAACTT TCAGAAGGGC CAGGTGACCC GAGACGGCTT CAAGCTGGTG ATGGCCTCCC | 240 |
|    | TGTACCACAT CTATGTGGCC CTGGAGGAGG AGATTGAGCG CAACAAGGAG AGCCCAGTCT | 300 |

- 27 -

TCGCCCCCTGT CTA CTTCCCA GAAGAGCTGC ACCGCAAGGC TGCCCTGGAG CAGGACCTGG 360  
CCTTCTGGTA CGGGCCCCGC TGGCAGGAGG TCATCCCCTA CACACCAGCC ATGCAGCGCT 420  
ATGTGAAGCG GCTCCACGAG GTGGGGCGCA CAGAGCCCGA GCTGCTGGTG GCCCACGCCT 480  
ACACCCGCTA CCTGGGTGAC CTGTCTGGGG GCCAGGTGCT CAAAAAGATT GCCCAGAAAG 540  
5 CCCTGGACCT GCCCAGCTCT GGCGAGGGCC TGGCCTTCTT CACCTTCCCC AACATTGCCA 600  
GTGCCACCAA GTTCAAGCAG CTCTACCGCT CCCGCATGAA CTCCCTGGAG ATGACTCCCC 660  
CAGTCAGGCA GAGGGTGATA GAAGAGGCCA AGACTGCGTT CCTGCTCAAC ATCCAGCTCT 720  
TTGAGGAGTT GCAGGAGCTG CTGACCATG ACACCAAGGA CCAGAGCCCC TCACGGGCAC 780  
CAGGGCTTCG CCAGCGGGCC AGCAACAAAG TGCAAGATTC TGCCCCCGTG GAGACTCCCA 840  
10 GAGGGAAGCC CCCACTCAAC ACCCGCTCCC AGGCTCCGCT TCTCCGATGG GTCCTTACAC 900  
TCAGCTTTCT GGTGGCGACA GTTGCTGTAG GGCTTTATGC CATGTGAATG CAGGCATGCT 960  
GGCTCCCAGG GCCATGAACT TTGTCCGTG GAAGGCCTTC TTTCTAGAGA GGAATTCTC 1020  
TTGGCTGGCT TCCTTACCGT GGGCACTGAA GGCTTTCAGG GCCTCCAGCC CTCTCACTGT 1080  
GTCCCTCTCT CTGGAAGGA GGAAGGAGCC TATGGCATCT TCCCCAACGA AAAGCACATC 1140  
15 CAGGCAATGG CCTAACTTC AGAGGGGGCG AAGGGGTCAG CCCTGCCCTT CAGCATCCTC 1200  
AGTTCCTGCA GCAGAGCCTG GAAGACACCC TAATGTGGCA GCTGTCTCAA ACCTCCAAAA 1260  
GCCCTGAGTT TCAAGTATCC TTGTTGACAC GGCCATGACC ACTTCCCCG TGGGCCATGG 1320  
CAATTTTAC ACAAACCTGA AAAGATGTTG TGTCTGTGT TTTTGTCTTA TTTTGTGTTG 1380  
AGCCACTCTG TTCCTGGCTC AGCCTCAAAT GCAGTATTTT TGTGTGTTC TGTGTTTTTT 1440  
20 ATAGCAGGGT TGGGGTGGTT TTTGAGCCAT GCGTGGGTGG GGAGGGAGGT GTTTAACGGC 1500  
ACTGTGGCCT TGGTCTAACT TTTGTGTGAA ATAATAACA ACATTGTCTG 1550

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 7 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

30 TGTGTCA

7

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 7 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double

- 28 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

TGAGTCA

7

What is claimed is:

1. A method of inhibiting sepsis-associated hypotension in a mammal, said method comprising identifying a mammal with or at risk of developing sepsis and administering to said mammal a compound which inhibits expression of inducible heme oxygenase (HO-1).
2. The method of claim 1, wherein said compound inhibits HO-1 transcription in a vascular cell of said mammal.
3. The method of claim 2, wherein said vascular cell is an aortic smooth muscle cell.
4. The method of claim 1, wherein said mammal is a human.
5. The method of claim 1, wherein said compound inhibits translation of HO-1 mRNA in a vascular cell of said mammal.
6. The method of claim 5, wherein said compound consists of a single stranded nucleic acid complementary to at least a portion of said HO-1 mRNA.
7. A method of inhibiting sepsis-associated hypotension in a mammal, said method comprising identifying a mammal with or at risk of developing sepsis and administering to said mammal a compound which inhibits the enzymatic activity of inducible heme oxygenase (HO-1).
8. The method of claim 7, wherein said compound is an antibody which binds to said HO-1.

- 30 -

9. A method for determining the ability of a candidate compound to inhibit HO-1 promoter activity, said method comprising the steps of:

- (a) providing DNA comprising a HO-1 promoter  
5 sequence linked to a reporter gene;
- (b) contacting said DNA with a candidate compound;  
and
- (c) determining the level of expression of said reporter gene, wherein a decrease in said level of  
10 expression in the presence of said compound, compared to the level of expression in the absence of said compound, is an indication of the ability of said candidate compound to inhibit HO-1 promoter activity.

10. A method for determining the ability of a  
15 candidate compound to inhibit expression of HO-1 in vascular smooth muscle cells, comprising

- (a) providing a vascular smooth muscle cell;
- (b) contacting said vascular smooth muscle  
cell with a candidate compound; and
- 20 (c) determining the amount of HO-1 expression in said vascular smooth muscle cell, wherein a decrease in said amount in the presence of said candidate compound compared to the amount in the absence of said candidate compound indicates that said candidate compound inhibits  
25 expression of HO-1 in vascular smooth muscle cells.

11. The method of claim 10, further comprising determining the amount of inducible nitric oxide synthase (iNOS) expression in said vascular smooth muscle cell, wherein a decrease in said amount of iNOS expression in  
30 the presence of said candidate compound compared to that in the absence of said candidate compound indicates that said candidate compound inhibits expression of both HO-1 and iNOS in vascular smooth muscle cells.



- 31 -

12. A transgenic non-human mammal the germ cells and somatic cells of which comprise a null mutation in a gene encoding HO-1.

13. The mammal of claim 12, wherein said mammal  
5 lacks enzymatically-active HO-1.

14. The mammal of claim 12, wherein an exon of said DNA is deleted.

15. The mammal of claim 12, wherein exon 3 of said DNA is deleted.

10 16. The mammal of claim 12, wherein said mammal is a rodent.



17. The mammal of claim 16, wherein said rodent is a mouse.

18. A mammalian cell line, the genomic DNA of  
15 which comprises a null mutation in DNA encoding HO-1, said cell line lacking the ability to synthesize enzymatically-active HO-1.



19. The cell line of claim 18, wherein said cell line is derived from a vascular smooth muscle cell.

20 20. The cell line of claim 19, wherein said cell line is immortalized.



1/4

**FIG. 1****FIG. 4**


2/4



3/4

**FIG. 5****FIG. 6**

4/4

**FIG. 7**

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US97/05440

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 39/395, 48/00; C12Q 1/02, 1/68; C12N 5/00, 15/00

US CL : 424/130.1 ; 435/6, 29, 325; 514/44; 800/2

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/130.1 ; 435/6, 29, 325; 514/44; 800/2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS; Chemical Abstracts

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SHIBAHARA, S et al. Functional Analysis of cDNAs for Two Types of Human Heme Oxygenase and Evidence for Their Separate Regulation. Journal of Biochemistry. 1993, Vol. 113, No. 2, pages 214-218, especially pages 216 and 218.	1-7 and 12-20
X	TRAKSHEL, G. M et al. Multiplicity fo Heme Oxygenase Isozymes. HO-1 and HO-2 are Different Molecular Species in Rat and Rabbit. Journal of Biological Chemistry. 15 January 1989, Vol. 264, No. 2, pages 1323-1328, especially pages 1327 and 1328.	7 and 9

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reasons (as specified)	*Z* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

15 JUNE 1997

Date of mailing of the international search report

10 JUL 1997

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DEBORAH CROUCH, PH.D.

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/05440

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	LAVROVSKY, Y. et al. Identification of Binding Sites for Transcription Factors NF-kappa-B and AP-2 in the Promoter Region fo the Human Heme Oxygenase 1 Gene. Proceeding of the National Academy of Sciences (USA), June 1994, Vol. 91, No. 13, pages 5987-5991, especially page 5990.	9